

2017-2021 Miami-Dade County Integrated HIV/AIDS Prevention and Care Plan

2018 IMPLEMENTATION PROGRESS

SECOND YEAR PLAN UPDATES

March 2018 – February 2019

Acronyms and Abbreviations Used in This Document

ACA	Affordable Care Act	NHAS	National HIV/AIDS Strategy
Acuity	Any of a number of co-occurring conditions or adherence issues contributing to an increased need for MCM attention	OHC	Oral Health Care, provided by the RWP
ADAP	AIDS Drug Assistance Program	OAHS	Outpatient/Ambulatory Health Services, provided by the RWP
AETC	AIDS Education and Training Center	OLTC	Joint RWP/FDOH-MDC Outreach Linkage to Care team
ARV	Antiretroviral Therapy	Part A/MAI	Part A and the Minority AIDS Initiative of the RWP
BSR	Behavioral Science Research Corp., (Admin/CQM Subrecipient)	PLWH	People Living With HIV
CQM	Clinical Quality Management Program at BSR	PLWHA	People Living With HIV/AIDS
CQMC	Ryan White Program CQM Committee	PrEP/nPEP	Pre-Exposure Prophylaxis /non-occupational Post-Exposure Prophylaxis
DIS	Disease Intervention Specialist at FDOH-MDC	PrEP WG	FDOH-MDC PrEP Work Group
FDOH-MDC	Florida Department of Health in Miami-Dade County	PRIM	Pre-Natal Immunology Clinic, w/in the University of Miami
FIMR	Fetal Infant Monitoring Review	RiC	Retention in Care
IDU	Injection Drug Use / Injection Drug User	Risk Factor	Self-reported mode of initial HIV/AIDS diagnosis
MAI	Minority AIDS Initiative, part of the RWP	RWP	Miami-Dade County Ryan White Program - Part A and MAI
MCM	RWP Medical Case Management or Medical Case Managers	STD	Sexually Transmitted Disease
MSM	Men Who Have Sex With Men	TTRA	Test and Treat / Rapid Access
		VL	Viral Load

Unless otherwise specified in the text of a strategy or activity, all strategies and activities are continuous progressive activities from January 1, 2017 to December 31, 2021.

**NHAS 2020 GOAL #1:
REDUCE NEW HIV INFECTIONS (2018 IMPLEMENTATION)**

PREVENTION

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source	
P1. By 2021, reduce new HIV infection rate by at least 25%, from 50.7 per 100,000 population in 2015 to 40.8 per 100,000 population in 2021, so that Miami-Dade County is no longer among the top three metropolitan areas with the highest incidence of new HIV infections.	PROGRESS TOWARD OBJECTIVE: Baseline 50.7 per 100,000 population in 2015, reduced to 40.8 per 100,000 in 2021.							
	P1.1 Increase access to and use of condoms by HIV positive persons and HIV-vulnerable populations, including but not limited to injection drug users (IDU), Trans-identified persons, gay and bisexual men.	P1.1a Increase the number of condom distribution sites. (added 2018 Q1)	FDOH-MDC and Partners	How many condoms were distributed to persons living with or at risk for HIV?	# of condoms distributed Distribution by location/Zip Code # of condoms distributed by Zip Code % of condoms distributed by Zip Code # of condoms distributed to ASOs, CBOs, treatment centers, HIV/IDU providers			Condom Distribution Coordinator Monthly reports
		P1.1b Develop an annual condom distribution map to identify new points of service (added Q3 2017)	FDOH-MDC and Partners					
		P1.1c Based on P1.1b data, recruit annually a new location or host an event to provide condom distribution services in the identified underserved area. (added 2018 Q1)	FDOH-MDC and Partners					
P1.1d Increase the availability and accessibility of condom distribution by 2.0 million a year		FDOH-MDC and Partners						

**NHAS 2020 GOAL #1:
REDUCE NEW HIV INFECTIONS (2018 IMPLEMENTATION)**

PREVENTION

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
<p>P1. By 2021, reduce new HIV infection rate by at least 25%, from 50.7 per 100,000 population in 2015 to 40.8 per 100,000 population in 2021, so that Miami-Dade County is no longer among the top three metropolitan areas with the highest incidence of new HIV infections.</p>	<p>P1.2 Implement STD and HIV testing to raise STD and HIV prevention awareness among HIV-vulnerable populations, including but not limited to IDU, Trans-identified persons, gay and bisexual men.</p>	<p>P1.2a Conduct HIV testing events in Miami Dade County.</p>	<p>FDOH-MDC and Partners</p>	<p>Was there an increase in HIV and STD testing among persons at risk?</p> <p>Was there an increase in the identification of HIV-negative persons at risk of HIV?</p> <p>Was there an increase in the number of persons living with HIV who are aware of their HIV or STD status?</p>	<p># of HIV tests # of STD tests</p> <p>% of tests that are among persons at risk for HIV</p> <p>% of tests stratified by priority target population (e.g. Hispanic MSM, IDU, Transgender persons, etc.)</p> <p>% of tests that are newly diagnosed with HIV.</p>		<p>1628 Testing Forms</p>
		<p>P1.2b Conduct STD testing events in Miami Dade County.</p>			<p>% of tests that are newly diagnosed with an STD</p>		

**NHAS 2020 GOAL #1:
REDUCE NEW HIV INFECTIONS (2018 IMPLEMENTATION)**

PREVENTION

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
<p>P1. By 2021, reduce new HIV infection rate by at least 25%, from 50.7 per 100,000 population in 2015 to 40.8 per 100,000 population in 2021, so that Miami-Dade County is no longer among the top three metropolitan areas with the highest incidence of new HIV infections.</p>	<p>P1.3 Implement HIV education to raise HIV prevention awareness among HIV-vulnerable populations, including but not limited to IDU, Trans-identified persons, gay and bisexual men.</p>	<p>P1.3a Partner with healthcare settings (e.g. hospitals, health centers, emergency departments), to increase the provision of routine HIV testing as part of medical care.</p>	<p>FDOH-MDC and Partners</p>	<p>NOTE: The annual County epidemiological profile of HIV cases breaks down by Zip Code. When the 2017 profile is fully available, a presentation will be scheduled.</p>			
		<p>P1.3b Increase the number of registered testing sites to ensure that HIV testing is more readily available and accessible</p>	<p>FDOH-MDC and Partners</p>				

**NHAS 2020 GOAL #1:
REDUCE NEW HIV INFECTIONS (2018 IMPLEMENTATION)**

PREVENTION

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
P2. Increase the number of individuals prescribed PrEP by at least 500% from the baseline 663 persons in 2016 to 3,978 persons by 2021.	P2.1 Increase availability of – and access to – PrEP/nPEP programs.	P2.1a Create a process for a PrEP external referral system (added Q3 2017).	FDOH-MDC and PrEP WG	Was there an increase in number of screenings of HIV-negative persons for PrEP eligibility? Was there an increase in PrEP-eligible persons?	% screened for PrEP % eligible for PrEP % linked to PrEP # counseled on PrEP # referred for PrEP		PrEPLink and HIV contracted provider monthly reports
		P2.1b Develop estimates of a PrEP cascade to inform prevention activities (began 01/01/18; dependent on P1.2a).	FDOH-MDC and PrEP WG	Were eligible persons linked to PrEP?	# prescribed PrEP		
		P2.1c Create a local directory of providers prescribing PrEP/nPEP, disseminate same on Part A and FDOH-MDC websites, and update annually thereafter.	FDOH-MDC and RWP				

**NHAS 2020 GOAL #1:
REDUCE NEW HIV INFECTIONS (2018 IMPLEMENTATION)**

PREVENTION

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
P3. Reduce the number of infants born with HIV in Miami-Dade County each year from three (3) to zero (0) by 2021.	P3.1 Increase number of OB/GYN healthcare providers engaging in HIV prevention activities with pregnant women.	P3.1b Conduct in-person educational sessions directed toward medical professionals who participate in the care of pregnant women with HIV, educating them about the requirements of Florida law and ensuring they are aware of community services available for women living with HIV and HIV exposed infants.	FDOH-MDC	Was there an increase in screening and active referral to prenatal HIV care among pregnant women living with diagnosed HIV?	% linked of post-partum women linked to family planning services / contraception services % of HIV positive pregnant women in HIV care % of presentation participants acknowledge an increase in awareness		Perinatal Coordinator Quarterly Reports
		P3.1c Conduct educational rounds with emergency rooms, urgent care centers, and classified high-risk delivery hospitals, to increase provider awareness of their responsibility to utilize the <i>High Risk Pregnancy Notification</i> and <i>Newborn Exposure Notification</i> forms and act on behalf of the pregnant women living with HIV and their HIV exposed babies.	FDOH-MDC				
		P3.1d Participate in an action-oriented community process to improve service systems and community resources for families.	FDOH-MDC and HBTF				

**NHAS 2020 GOAL #1:
REDUCE NEW HIV INFECTIONS (2018 IMPLEMENTATION)**

PREVENTION

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
P3. Reduce the number of infants born with HIV in Miami-Dade County each year from three (3) to zero (0) by 2021.	P3.2 Conduct targeted public information campaigns toward pregnant women at risk of HIV, to have access to OB/GYN providers, HIV prevention materials and information on community services for women with HIV/AIDS.	P3.2a Conduct community outreach and promote information campaigns towards women of child-bearing age living with HIV.	FDOH-MDC	How many agencies are providing post-partum family planning services to women living with HIV?	# of agencies		
		P3.2b Create linkage services assuring at least 90% of post-partum women living with HIV have access to contraceptive/ family planning and preconception care services after delivery (no baseline).	FDOH-MDC and PRIM				

NHAS 2020 GOAL #2:

INCREASE ACCESS TO CARE AND IMPROVE HEALTH OUTCOMES FOR PEOPLE LIVING WITH HIV OR AIDS

LINKAGE TO CARE

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
<p>L 1. Increase the percentage of PLWH linked to medical care [at least one (1) outpatient medical care visit, as documented by a billed OAHS encounter, or a CD4 or VL test] within 30 days post-diagnosis to at least 85% by 2021. [55% based on 90 days post-diagnosis]</p>	<p>L 1.1 Improve existing FDOH-Part A diagnosis-to-linkage client management process.</p> <p>Note: During CY 2018, TTRA became the de facto “best practice” for new-to-care PLWH.</p>	<p>PROGRESS TOWARD OBJECTIVE: Baseline post-diagnosis linkage rate of 55% within 90 days of initial diagnosis in 2015, increased to 85% by 2021 (starting Second Year: Linkage within 30 days).</p>					
		<p>L1.1a Evaluate the FDOH-MDC / RWP Test and Treat/Rapid Access (TTRA) process and make improvements wherever possible.</p>	<p>FDOH-MDC and Partners</p> <p>RWP CQM</p>	<p>Is there an increase in the percentage of newly-diagnosed PLWH linked to care within 30 days of diagnosis? [Baseline 2018 data]</p> <p>What percentage of TTRA clients are enrolled in ADAP within 30 days of the first OAHS visit?</p>	<p># newly-diagnosed PLWH by subrecipient and total [FDOH-MDC]</p> <p># TTRA PLWH linked to RWP care</p> <p>% newly-diagnosed PLWH linked to RWP care through TTRA within 30 days of diagnosis</p>		
		<p>L1.1c Identify percentage of newly-diagnosed TTRA clients who access ARVs within 30 days of first OAHS visit.</p>	<p>FDOH-MDC</p> <p>RWP CQM</p>		<p># TTRA clients enrolled in ADAP within 30 days of the first OAHS visit</p>		
		<p>L1.1d FDOH-MDC will hold monthly 501 Update training, including RWP linkage and outreach updates, for counseling and testing site coordinators.</p>	<p>FDOH-MDC</p> <p>RWP CQM</p>		<p>% TTRA clients enrolled in ADAP within 30 days of the first OAHS visit</p>		

NHAS 2020 GOAL #2:

INCREASE ACCESS TO CARE AND IMPROVE HEALTH OUTCOMES FOR PEOPLE LIVING WITH HIV OR AIDS

LINKAGE TO CARE

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
<p>L1. Increase the percentage of PLWHA linked to medical care [at least one (1) outpatient medical care visit, as documented by a billed OMC encounter, or a CD4 or VL test] within 30 days post-diagnosis from 55% in 2015 to at least 85% by 2021.</p>	<p>L1.2 Provide Partner Services to identified HIV+ individuals, allowing for the notification, screening and referral to appropriate services for partners of newly-diagnosed PLWHA.</p>	<p>L1.2a Increased participation in HIV partner services among persons with diagnosed HIV (baseline: 68% in 2015)</p>	<p>FDOH-MDC DIS</p>	<p>Was there an increase in notification in HIV testing of partners identified through HIV partner services?</p> <p>What is the impact of Partner Services on engagement, testing and linkage of PLWHA who are partners of persons diagnosed with HIV?</p>	<p>Of all named, notifiable partners identified through HIV partner services, the percentage notified for HIV partner services</p> <p>% named, notifiable partners that were tested for HIV</p> <p># of newly diagnosed PLWHA with stable partners, who are potentially eligible for Partner Services, at beginning of baseline evaluation period, and at 6 and 12 months thereafter</p> <p>% of partners of newly diagnosed PLWHA who are engaged in Partner Services, at beginning of baseline period and at 6 and 12 months thereafter</p>		<p>Partner Service Reports</p>
		<p>L1.2b Increased notification and HIV testing of partners identified through HIV partner services. (baseline: 48% in 2015)</p>	<p>FDOH-MDC DIS</p>				
		<p>L1.2c Explore creating a video direct observational therapy system for partners (began 01/01/18).</p>	<p>FDOH-MDC</p>				

NHAS 2020 GOAL #2:

INCREASE ACCESS TO CARE AND IMPROVE HEALTH OUTCOMES FOR PEOPLE LIVING WITH HIV OR AIDS

LINKAGE TO CARE

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
<p>L1. Increase the percentage of PLWHA linked to medical care [at least one (1) outpatient medical care visit, as documented by a billed OMC encounter, or a CD4 or VL test] within 30 days post-diagnosis from 55% in 2015 to at least 85% by 2021.</p>	<p>L1.3 Identify and link to medical care at least 25% of the newly-diagnosed HIV+ persons identified through the FDOH-MDC Data To Care (DTC) initiative.</p>	<p>L1.3a Provide linkage to, re-engagement in, and retention in RWP HIV medical services using DTC activities.</p>	<p>FDOH-MDC RWP CQM</p>	<p>Was there an increase in linkage of persons to HIV medical care attributable to DTC?</p>	<p>% not in care % reengaged in care</p>		

NHAS 2020 GOAL #2:

INCREASE ACCESS TO CARE AND IMPROVE HEALTH OUTCOMES FOR PEOPLE LIVING WITH HIV OR AIDS

RETENTION IN CARE

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
R1. Increase the percentage of RWP clients who had at least two (2) instances of either (a) RWP OAHS visits, or (b) CD4/VL lab tests (as proxy for medical care outside the RWP), at least 90 days apart within a 12 month period, from 60% in 2015 to at least 90% by 2021.	R1.1 Identify RWP client target populations who are at greatest risk for dropping out of care. NOTE: All HRSA demographics will be reviewed by the Strategic Planning Committee.	R1.1a Identify RWP client demographic characteristics (ethnicity, gender, age) and risk factor associated with low Retention in Care (RiC) rates and track RiC rates by demographic and risk factor groups across and within subrecipients.	RWP CQM	What are the RiC rates by demographic characteristics? What are the RiC rates by risk factors? What are the RiC rates by co-occurring conditions/acuties?	# of RWP clients receiving MCM and OAHS at the beginning of evaluation period % of RWP clients RiC by subrecipient at 6 and 12 months thereafter % of RWP clients RiC by demographic characteristics		SDIS
		R1.1b Develop assessments of acuties (e.g. substance use, mental illness, incidence of missed appointments or other non-adherence) associated with dropping out of care; track and refine measurements in subsequent years.	RWP CQM	What are the potential problem areas to remediate? What are the best practices to replicate?	% of RWP clients RiC by risk factors % of RWP clients RiC by co-occurring conditions/acuties		
		R1.1c Identify RWP subrecipients with highest and lowest RiC rates, identify potential problem areas to remediate and best practices to replicate.	RWP CQM	What is the impact of replication of a subrecipient best practice for improving RiC rates on actual RiC?			

NHAS 2020 GOAL #2:

INCREASE ACCESS TO CARE AND IMPROVE HEALTH OUTCOMES FOR PEOPLE LIVING WITH HIV OR AIDS

RETENTION IN CARE

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
R1. Increase the percentage of RWP MCM clients who had at least two (2) instances of either (a) RWP OAHS visits, or (b) CD4/VL lab tests (as proxy for medical care outside the RWP), at least 90 days apart within a 12 month period, from 60% in 2015 to at least 90% by 2021.	R1.4 Provide continuous improvements in MCM and OAHS that meet the needs and identified vulnerabilities of PLWH in care.	R1.4a Monitor quality and appropriateness of MCM and OAHS clinical care through client satisfaction surveys, targeted reviews of SDIS data and clinical record reviews, provide technical assistance to assist subrecipients in self-correction (e.g., on site CQM TA, AETC, etc.). Provide subrecipient-based data to the Recipient and CQMC to make recommendations in service delivery processes.	RWP CQM CQMC	Are there significant relationships between subrecipients' client satisfaction scores for MCM and OAHS service delivery and clinical outcomes (RiC, VL suppression) for PLWH clients served by those subrecipients?	# of OAHS, OHC and MCM subrecipients. # of OAHS subrecipients who received an SF-SE AETC Record Review. # of OAHS subrecipients with record review data forwarded to OMB.		SDIS CQM client surveys CQM record reviews
		R1.4c Review and update RWP Service Delivery Guidelines for RWP OAHS, OHC, and MCM services annually.	RWP CQM OMB	Are the RWP OAHS, OHC, and MCM subrecipients fully engaged in review and response to Service Delivery Guidelines updates?	# of MCM and OHC subrecipients identified with service delivery issues resulting in record reviews.		
		R1.4d Conduct record reviews of RWP OAHS, OHC, and MCM subrecipients to ensure adherence to PHS and RWP Service Delivery Guidelines.	RWP CQM SF-SE AETC OMB	What areas are identified as needing improvement?	# of MCM and OHC subrecipients with record review data forwarded to OMB.		
		R1.4e Provide Peer capacity-building activities that address RiC.	RWP CQM	How many RWP subrecipients received record reviews?			

NHAS 2020 GOAL #2:

INCREASE ACCESS TO CARE AND IMPROVE HEALTH OUTCOMES FOR PEOPLE LIVING WITH HIV OR AIDS

RETENTION IN CARE

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
<p>R1. Increase the percentage of RWP MCM clients who had at least two (2) instances of either (a) RWP OAHS visits, or (b) CD4/VL lab tests (as proxy for medical care outside the RWP), at least 90 days apart within a 12 month period, from 60% in 2015 to at least 90% by 2021.</p>	<p>R1.5 Track and assess the quality of OAHS care provided to RWP clients who transition to ACA-provided outpatient medical care.</p>	<p>R1.5a Revise and streamline ACA enrollment and re-enrollment processes to increase the number of ACA-eligible RWP clients transitioned to ACA from 50% (2015 baseline) to 70%.</p>	<p>RWP CQM</p>	<p>Is care delivered through the ACA doing an effective job of maintaining and improving clinical outcomes for RWP clients in care?</p>	<p># of RWP clients determined eligible for ACA enrollment, as of annual enrollment period.</p>		<p>SDIS</p>
		<p>R1.5b Require all RWP clients enrolled in ACA and receiving RWP MCM to report VL levels at a minimum once every six (6) months, in order to assess and track clinical health outcomes of RWP clients receiving OAHS through an ACA Marketplace insurance plan.</p>	<p>RWP CQM</p>	<p>Is client satisfaction higher among ACA enrollees vs. RWP? Are VL suppression rates higher among ACA enrollees vs. RWP?</p>	<p># of RWP clients enrolled in ACA plans as of annual enrollment period. % of ACA-eligible PLWH enrolled.</p>		
		<p>R1.5c Compare rates of missing VL data and VL suppression rates among PLWH treated through the ACA with missing VL data and VL suppression rates among PLWH receiving OAHS through the RWP to identify disparities.</p>	<p>RWP CQM</p>	<p>Are differences in clinical outcomes shown between ACA-enrolled clients and RWP clients related to:</p>	<p>Client outcome data (VL suppression, RiC) for clients eligible for ACA, vs. clients enrolled in ACA, vs. RWP clients not eligible for ACA.</p>		
		<p>R1.5d Increase the percentage of clients transitioned from RWP-funded OAHS to ACA medical care who are retained in ACA-provided medical care for two (2) years after enrollment from 60% enrolled in 2015 and continuously enrolled in 2017 to 75% enrolled in 2019 and continuously enrolled in 2021. (Using VL data as proxy for ACA OAHS)</p>	<p>RWP CQM</p>	<p>a) differences in the characteristics of PLWH who can or cannot enroll in ACA, or b) differences in the levels of care provided through ACA coverage vs. RWP care?</p>			

NHAS 2020 GOAL #2:

INCREASE ACCESS TO CARE AND IMPROVE HEALTH OUTCOMES FOR PEOPLE LIVING WITH HIV OR AIDS

RETENTION IN CARE

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
R2. Increase the proportion of "lost to care" RWP MCM clients who are re-linked to care, from 40% in 2017 to 60% by 2021.	R2.1 RWP MCM subrecipients will partner with FDOH-MDC surveillance to detect clients in danger of being lost to care, update contact information on vulnerable RWP clients, and use both FDOH and RWP outreach specialists to re-link clients in care.	R2.1a The joint FDOH/RWP OLTC Team will determine criteria for "truly lost to care" case closures in MCM subrecipients and develop an "early warning system" to identify RWP clients at risk for being lost to care.	FDOH-MDC RWP CQM	Does the "early warning system" improve the RiC rates for PLWH in MCM care? Are there key client groups (demographic, acuity, co-occurring conditions) that are more likely to show "early warning indicators" for being lost to care?	# of PLWH at each MCM subrecipient provider at beginning of evaluation period. % of PLWH with certifiably closed cases at each MCM subrecipient site at 6 and 12 months.		SDIS
		R2.1b FDOH and RWP will develop data-sharing protocols and feedback mechanisms to provide updated contact information to RWP on clients who are flagged by the "early warning system" as at risk for being lost to care, as well as provide case closure data to FDOH for clients with 6, 9 and 12 months since the most recent VL measurement or on-site RWP OAHS contact (began 01/01/18).	FDOH-MDC RWP CQM	After CQM intervention, are there demonstrable improvements in re-linkage rates among subrecipients with low re-linkage rates?	% of PLWH identified by the early warning system as RiC risks, at each MCM subrecipient site. % of PLWH identified by the early warning system who are lost to care, at each MCM subrecipient site, at 6 and 12 months.		
		R2.1c Identify RWP MCM subrecipients with lowest and highest re-linkage rates, and determine best practices that may be disseminated and adopted within the RWP system.	FDOH-MDC RWP CQM	After "best practice" intervention, is there improvement in RiC and re-linkage?			

NHAS 2020 GOAL #2:

INCREASE ACCESS TO CARE AND IMPROVE HEALTH OUTCOMES FOR PEOPLE LIVING WITH HIV OR AIDS

VIRAL LOAD SUPPRESSION

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
<p>V1. Increase the percentage of people living with HIV in the EMA who are virally suppressed (<200 copies/mL) from 67% in 2015 to at least 80% by 2021.</p>	<p>V1.3 Expand role of RWP MCM and OAHS subrecipients in detecting persistent unsuppressed viral loads (VL) and initiate appropriate responses.</p>	<p>V1.3a On a monthly basis, detect PLWH with persistent unsuppressed VL over two consecutive semi-annual measurements and notify RWP MCM and OAHS subrecipients to enable their targeted response. Monitor improvement in VL suppression levels to ensure efficacy of subrecipient response.</p> <p>Note: OAHS data will be limited to RWP OAHS subrecipients only.</p>	<p>RWP CQM CQMC</p>	<p>Which MCM and OAHS subrecipients have the highest proportion of clients with new or continued persistent unsuppressed VL?</p> <p>Can alerting RWP subrecipients about clients with persistent unsuppressed VL produce internal responses to address client VL issues?</p> <p>What type of subrecipient service – MCM or OAHS – is more effective in addressing persistent unsuppressed VL when informed?</p>	<p>% of clients served by MCM and OAHS subrecipients who show persistent unsuppressed VL.</p> <p>% of identified clients with persistent unsuppressed VL among MCM and OAHS subrecipients, and whose VL levels show improvement at next semi-annual measurement.</p>		<p align="center">SDIS</p>

**NHAS 2020 GOAL #3:
REDUCE HIV-RELATED HEALTH DISPARITIES AND HEALTH INEQUITIES (2018 IMPLEMENTATION)**

DISPARITIES IN RETENTION IN CARE

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
<p>DR1. By 2021, increase the percentage of RWP MCM clients in three (3) key disparate minority health groups who had at least two (2) instances of either (a) RWP OAHS visits, or (b) CD4/VL lab tests (as proxy for medical care outside the RWP), at least 90 days apart within a 12 month period, from 60% in 2015 to at least 90% by 2021, to match RWP total levels.</p>	<p>DR1.1 Identify retention in care (RiC) vulnerabilities of Black/African American male (B/AAM) clients within the RWP network of care, and address them with specific interventions: Increase RiC among B/AAM from 53% in 2015 to 90% by 2021.</p>	<p>DR1.1a Determine (1) best practices of RWP MCM subrecipients with higher than average PLWH RiC rates for B/AAM; and (2) risk factors and acuities contributing to low B/AAM PLWH RiC rates.</p>	RWP CQM	What are the risk-factor/acuity-related RiC rates for B/AAM?	% RiC for B/AAM by individual MCM subrecipients.		SDIS
		<p>DR1.1b Identify MAI program RiC initiatives of RWP MCM subrecipients directed toward B/AAM, and track RiC rates annually.</p>	RWP CQM	What contributes to B/AAM RiC successes among MCM subrecipients? Are these "best practices" that may be replicated?	% RiC for B/AAM in care with MCM subrecipients by risk factors/acuities.		
		<p>DR1.1c Evaluate best practices and MAI program RiC initiatives directed toward B/AAM for replicability, select one or more for replication pilot-testing and evaluation to improve RiC rates for B/AAM PLWH in RWP-funded MCM subrecipients.</p> <p>Note: System-wide replication may be recommended but cannot be mandated.</p>	RWP CQM	What MAI-funded program initiatives have positive impacts on RiC for B/AAM? What is the impact of replication of a subrecipient best practice or MAI initiative on RiC rates for B/AAM?	% RiC for B/AAM clients of MCM subrecipients with identified "best practices" and MAI-funded program initiatives. % RiC for B/AAM clients of subrecipients where "best practices" and/or MAI-funded initiatives have been replicated.		

**NHAS 2020 GOAL #3:
REDUCE HIV-RELATED HEALTH DISPARITIES AND HEALTH INEQUITIES (2018 IMPLEMENTATION)**

DISPARITIES IN RETENTION IN CARE

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
<p>DR1. By 2021, increase the percentage of RWP MCM clients in three (3) key disparate minority health groups who had at least two (2) instances of either (a) RWP OAHS visits, or (b) CD4/VL lab tests (as proxy for medical care outside the RWP), at least 90 days apart within a 12 month period, from 60% in 2015 to at least 90% by 2021, to match RWP total levels.</p>	<p>DR1.2 Identify retention in care (RiC) vulnerabilities of Black/African American female (B/AAF) clients within the RWP network of care, and address them with specific interventions: Increase RiC rates for B/AAF from 52% in 2015 to 90% by 2021.</p>	<p>DR1.2a Determine (1) best practices of RWP MCM subrecipients with higher than average PLWH RiC rates for B/AAF; and (2) risk factors and acuities contributing to low B/AAF PLWH RiC rates.</p>	RWP CQM	What are the risk-factor/acuity-related RiC rates for B/AAF?	% RiC for B/AAF by individual MCM subrecipients.		SDIS
		<p>DR1.2b Identify MAI program RiC initiatives of RWP MCM subrecipients directed toward B/AAF, and track RiC rates annually.</p>	RWP CQM	What contributes to B/AAF RiC successes among MCM subrecipients? Are these "best practices" that may be replicated?	% RiC for B/AAF in care with MCM subrecipients by risk factors/acuities.		
		<p>DR1.2c Evaluate best practices and MAI program RiC initiatives directed toward B/AAF for replicability, select one or more for replication pilot-testing and evaluation to improve RiC rates for B/AAF PLWH in RWP-funded MCM subrecipients.</p> <p>Note: System-wide replication may be recommended but cannot be mandated.</p>	RWP CQM	What MAI-funded program initiatives have positive impacts on RiC for B/AAF? What is the impact of replication of a subrecipient best practice or MAI initiative on RiC rates for B/AAF?	% RiC for B/AAF clients of MCM subrecipients with identified "best practices" and MAI-funded program initiatives. % RiC for B/AAF clients of subrecipients where "best practices" and/or MAI-funded initiatives have been replicated.		

**NHAS 2020 GOAL #3:
REDUCE HIV-RELATED HEALTH DISPARITIES AND HEALTH INEQUITIES (2018 IMPLEMENTATION)**

DISPARITIES IN RETENTION IN CARE

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
<p>DR1. By 2021, increase the percentage of RWP MCM clients in three (3) key disparate minority health groups who had at least two (2) instances of either (a) RWP OAHS visits, or (b) CD4/VL lab tests (as proxy for medical care outside the RWP), at least 90 days apart within a 12 month period, from 60% in 2015 to at least 90% by 2021, to match RWP total levels.</p>	<p>DR1.3 Identify retention in care (RiC) vulnerabilities of Hispanic MSM (HMSM) clients within the RWP network of care, and address them with specific interventions: Increase RiC rates among HMSM from 69%* in 2015 to 90% by 2021.</p> <p><i>*Based on FDOH 2015 Epi data.</i></p>	<p>DR1.3a Determine (1) best practices of RWP MCM subrecipients with higher than average PLWH RiC rates for HMSM; and (2) risk factors and acuities contributing to low HMSM PLWH RiC rates.</p>	<p>RWP CQM CQMC</p>	<p>What are the risk-factor/acuity-related RiC rates for HMSM?</p> <p>What contributes to HMSM RiC successes among MCM subrecipients? Are these "best practices" that may be replicated?</p> <p>What MAI-funded program initiatives have positive impacts on RiC for HMSM?</p>	<p>% RiC for HMSM by individual MCM subrecipients.</p> <p>% RiC for HMSM in care with MCM subrecipients by risk factors/acuities.</p> <p>% RiC for HMSM clients of MCM subrecipients with identified "best practices" and MAI-funded program initiatives.</p>		SDIS
		<p>DR1.3b Identify MAI program RiC initiatives of RWP MCM subrecipients directed toward HMSM, and track RiC rates annually.</p>	<p>RWP CQM CQMC</p>	<p>What is the impact of replication of a subrecipient best practice or MAI initiative on RiC rates for HMSM?</p>	<p>% RiC for HMSM clients of subrecipients where "best practices" and/or MAI-funded initiatives have been replicated.</p>		
		<p>DR1.3c Evaluate best practices and MAI program RiC initiatives directed toward HMSM for replicability, select one or more for replication pilot-testing and evaluation to improve RiC rates for HMSM PLWH in RWP-funded MCM subrecipients.</p> <p>Note: System-wide replication may be recommended but cannot be mandated.</p>	<p>RWP CQM CQMC</p>				

**NHAS 2020 GOAL #3:
REDUCE HIV-RELATED HEALTH DISPARITIES AND HEALTH INEQUITIES (2018 IMPLEMENTATION)**

DISPARITIES IN TREATMENT OUTCOMES (SUPPRESSED / UNDETECTABLE VL LEVELS)

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
DV1. By 2021, increase the percentage of RWP MCM clients in three (3) key disparate minority health groups receiving who have suppressed viral loads, to 80% by 2021, to match overall PLWH levels.	DV1.1 Identify VL suppression vulnerabilities associated with Black/African American male (B/AAM) clients within the RWP network of care, and address them with specific interventions: Increase B/AAM VL suppression levels from 57% suppressed VL in 2015 to 80% by 2021.	DV1.1a Determine (1) best practices of RWP MCM subrecipients with higher than average PLWH VL suppression rates for B/AAM; and (2) risk factors and acuities contributing to low B/AAM PLWH VL suppression rates.	RWP CQM	What are the risk-factor/acuity-related VL suppression rates for B/AAM? What contributes to B/AAM VL suppression successes among MCM subrecipients? Are these "best practices" that may be replicated?	% suppressed VL for B/AAM by individual MCM subrecipients. % suppressed VL for B/AAM in care with MCM subrecipients by risk factors/acuities.		SDIS
		DV1.1b Identify MAI program VL suppression initiatives of RWP MCM subrecipients directed toward B/AAM, and track VL suppression rates annually.	RWP CQM	What MAI-funded program initiatives have positive impacts on VL suppression for B/AAM?	% suppressed VL for B/AAM clients of MCM subrecipients with identified "best practices" and MAI-funded program initiatives.		
		DV1.1c Evaluate best practices and MAI program VL suppression initiatives directed toward B/AAM for replicability, select one or more for replication pilot-testing and evaluation to improve VL suppression rates for B/AAM PLWH in RWP-funded MCM subrecipients. Note: System-wide replication may be recommended but cannot be mandated.	RWP CQM	What is the impact of replication of a subrecipient best practice or MAI initiative on VL suppression rates for B/AAM?	% suppressed VL for B/AAM clients of subrecipients where "best practices" and/or MAI-funded initiatives have been replicated.		

**NHAS 2020 GOAL #3:
REDUCE HIV-RELATED HEALTH DISPARITIES AND HEALTH INEQUITIES (2018 IMPLEMENTATION)**

DISPARITIES IN TREATMENT OUTCOMES (SUPPRESSED / UNDETECTABLE VL LEVELS)

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
DV1. By 2021, increase the percentage of RWP MCM clients in three (3) key disparate minority health groups receiving who have suppressed viral loads, to 80% by 2021, to match overall PLWH levels.	DV1.2 Identify VL suppression vulnerabilities associated with Black/African American female (B/AAF) clients within the RWP network of care, and address them with specific interventions: Increase B/AAF VL suppression levels from 60% suppressed VL in 2015 to 80% by 2021.	DV1.2a Determine (1) best practices of RWP MCM subrecipients with higher than average PLWH VL suppression rates for B/AAF; and (2) risk factors and acuities contributing to low B/AAF PLWH VL suppression rates.	RWP CQM CQMC	What are the risk-factor/acuity-related VL suppression rates for B/AAF? What contributes to B/AAF VL suppression successes among MCM subrecipients? Are these "best practices" that may be replicated? What MAI-funded program initiatives have positive impacts on VL suppression for B/AAF? What is the impact of replication of a subrecipient best practice or MAI initiative on VL suppression rates for B/AAF?	% suppressed VL for B/AAF by individual MCM subrecipients. % suppressed VL for B/AAF in care with MCM subrecipients by risk factors/acuities. % suppressed VL for B/AAF clients of MCM subrecipients with identified "best practices" and MAI-funded program initiatives. % suppressed VL for B/AAF clients of subrecipients where "best practices" and/or MAI-funded initiatives have been replicated.		SDIS
		DV1.2b Identify MAI program VL suppression initiatives of RWP MCM subrecipients directed toward B/AAF, and track VL suppression rates annually.	RWP CQM CQMC				
		DV1.2c Evaluate best practices and MAI program VL suppression initiatives directed toward B/AAF for replicability, select one or more for replication pilot-testing and evaluation to improve VL suppression rates for B/AAF PLWH in RWP-funded MCM subrecipients. Note: System-wide replication may be recommended but cannot be mandated.	RWP CQM CQMC				

**NHAS 2020 GOAL #3:
REDUCE HIV-RELATED HEALTH DISPARITIES AND HEALTH INEQUITIES (2018 IMPLEMENTATION)**

DISPARITIES IN TREATMENT OUTCOMES (SUPPRESSED / UNDETECTABLE VL LEVELS)

Objectives	Strategies	Activities	Responsible Entities	Evaluation Questions	Outputs	YTD Data	Data Source
DV1. By 2021, increase the percentage of RWP MCM clients in three (3) key disparate minority health groups receiving who have suppressed viral loads, to 80% by 2021, to match overall PLWH levels.	DV1.3 Identify VL suppression vulnerabilities associated with Haitian clients within the RWP network of care, and address them with specific interventions: Increase Haitian VL suppression levels from 68% suppressed VL in 2015 to 80% by 2021.	DV1.3a Determine (1) best practices of RWP MCM subrecipients with higher than average PLWH VL suppression rates for Haitians; and (2) risk factors and acuities contributing to low Haitian PLWH VL suppression rates.	RWP CQM CQMC	What are the risk-factor/acuity-related VL suppression rates for Haitians? What contributes to Haitian VL suppression successes among MCM subrecipients? Are these "best practices" that may be replicated?	% suppressed VL for Haitian PLWHA by individual MCM subrecipients. % suppressed VL for Haitians in care with MCM subrecipients by risk factors/acuities. % suppressed VL for Haitian clients of MCM subrecipients with identified "best practices" and MAI-funded program initiatives.		SDIS
		DV1.3b Identify MAI program VL suppression initiatives of RWP MCM subrecipients directed toward Haitians, and track VL suppression rates annually.	RWP CQM CQMC	What MAI-funded program initiatives have positive impacts on VL suppression for Haitian clients? What is the impact of replication of a subrecipient best practice or MAI initiative on VL suppression rates for Haitian clients?	% suppressed VL for Haitian clients of subrecipients where "best practices" and/or MAI-funded initiatives have been replicated.		
		DV1.3c Evaluate best practices and MAI program VL suppression initiatives directed toward Haitians for replicability, select one or more for replication pilot-testing and evaluation to improve VL suppression rates for Haitian PLWH in RWP-funded MCM subrecipients. Note: System-wide replication may be recommended but cannot be mandated.	RWP CQM CQMC				

Objectives, Strategies, and Activities completed and/or outside the scope of the Responsible Entities:

- **L 1.1b** Create a comprehensive outreach linkage resource directory, mapping outreach service providers by area within Miami-Dade County, and ensuring that at least one outreach provider in each area provides outreach/linkage services outside normal business hours and on weekends.
- **L1.1e** Guarantee flexible after-hours and weekend availability at RWP MCM, OAHS and OHC subrecipients, ensuring that at least one (1) MCM subrecipient, one (1) OAHS subrecipient and one (1) OHC subrecipient will offer after-hours service five (5) weeknights per week and will offer four (4) hours of service availability one (1) weekend day, to reduce or eliminate barriers to RWP linkage once a PLWH has been diagnosed at FDOH-MDC site.
- **R1.2a** Identify service improvement opportunities to improve RiC rates, to R1.1.
- **OLD R1.2b** With the CQMC, identify one replicable best MCM program practice for RiC, replicate it as a pilot program and evaluate its impact in another MCM agency.
- **R1.2b4** Monitor and improve clinical proficiency among RWP MCM, outreach workers, and peers, through online and group proficiency development and clinical training.
- **R1.3** Enhance the RWP Client-Centered Care quality management protocols for (1) reducing lagtime to first or repeat appointments; (2) reducing wait time in lobby for appointments; and (3) reducing hold time reaching live help by telephone, for Ryan White Program OAHS, OHC and MCM subrecipients.
 - **R1.3a** Assess measurement systems for measuring lagtime, wait time and hold time among RWP OAHS, OHC and MCM subrecipients, and track subrecipients thereafter.
 - **R1.3b** Assess RiC rates of subrecipient MCM and OAHS providers based on lagtime wait time. and hold time.
 - **R1.3c** Reduce average lagtime, average wait time and average hold time by 25%, relative to standards updated in R1.2c and tracked in R1.3a.
 - **R1.3d** In conjunction with the CQMC, identify at least one (1) program-based lagtime, wait time or hold time best practice or Quality Improvement opportunity resulting in improved RiC.

Per Recipient, this will no longer be tracked since Client Satisfaction Survey indicates high satisfaction and there is no indication of change in RiC or VL suppression based on lagtime, wait time or hold time. RWP subrecipients are contractually required to meet MCM, AOHS, and OHC appointment setting standards defined by the Recipient.

- **P1.3c** Rebrand the STD clinic name to promote sexual health and wellness (began 01/01/18).
- **P1.4** Achieve a more coordinated local response to the HIV epidemic.
 - **P1.4a** Publish approved updates to the Integrated Plan (IP) on Test Miami and RWP websites, promoting awareness.
- **P2.1d** By December 31, 2019 (Third Year), facilitate establishment of new PrEP/nPEP medical providers.
- **P2.1e** Establish a “PrEP Ambassador” program.
- **OLD P2.1a** Conduct an annual outbound direct mail and email awareness campaign to inform 600+ Miami-Dade OB/GYN physicians and physician assistants of (1) state law 64D-3.042 and (2) the local protocol of High Risk Pregnancy Notification and Newborn Exposure Notification, (3) the Protect Yourself, Protect Your Baby (PYPYB) campaign, and (4) to encourage awareness of their responsibility to act on behalf of the HIV infected pregnant women with HIV in their care. The mail-out will include bounce-back cards to enumerate the level of cooperation and the number of pregnant women with HIV infected pregnant women in care, and to allow providers to request more information.

- **OLD P2.2** Increase the engagement of OB/GYN providers and the community in prevention of perinatally acquired HIV.
 - **OLD P2.2a** Participate in FIMR-HIV meetings to review all information gathered on perinatal transmission cases, identify system issues, and make recommendations for improvement.
 - **OLD P2.2b** Participate in Community Action Team meetings to initiate systems change based on FIMR-HIV recommendations for improvement.
 -
 - **OLD P3.2a** Disseminate educational materials, promoting the PYPYB awareness campaign, encouraging women of childbearing age who are pregnant or considering pregnancy to get tested for HIV (see also P2.1a).
 - **OLD P3.2b** Partner with at obstetric healthcare providers, who agree to show the PYPYB DVD in patient waiting rooms (no baseline).

- **D1.** Increase the percentage of PLWH who know their serostatus from 86% in 2015 to at least 90% by 2021.
 - **D1.1** Partner with healthcare settings (e.g. hospitals, health centers, emergency departments), to increase the provision of routine HIV testing as part of medical care. **NOTE:** 30% of emergency visits (224,989 / 754,362) are receiving an HIV test and will increase to 41% (316,845 / 754,362) by the end of 2018. The three largest emergency rooms in Miami-Dade County will or are implementing routinized HIV testing.
 - **D1.2** Intensify HIV efforts in high impact areas.

- **V1.1** Provide immediate access to antiretroviral therapy through timely linkage to medical care, expanding the FDOH-MDC Test and Treat model of care.
- **V1.2** Enhance support for FDOH-MDC treatment adherence (FDOH-MDC initiatives).